

Sum/difference - to - product formulae

Na prezentacji omówimy wzory z ostatniej lekcji i pokażemy jak je stosować przy rozwiązywaniu równań trygonometrycznych.

When we solve an equation like:

$$x^2 - 4x - 12 = 0 \quad (1)$$

what we want to do is to turn it (if possible) into:

$$(x - 6)(x + 2) = 0 \quad (2)$$

because this immediately gives us the solutions $x = 6$ or $x = -2$.

When we solve an equation like:

$$x^2 - 4x - 12 = 0 \quad (1)$$

what we want to do is to turn it (if possible) into:

$$(x - 6)(x + 2) = 0 \quad (2)$$

because this immediately gives us the solutions $x = 6$ or $x = -2$.

We turned equation (1) where we add terms to an equation (2) where we multiply terms. This is often useful when solving equations.

We want to do something similar to trigonometric functions. Suppose we have an equation:

$$\sin A + \cos B = 0$$

can we somehow turn this sum into a product?

We want to do something similar to trigonometric functions. Suppose we have an equation:

$$\sin A + \cos B = 0$$

can we somehow turn this sum into a product?

The answer is of course yes and we will see how to do this on the next few slides.

Recall that we already have the following four formulae:

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \sin \beta \cos \alpha$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$.

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$. This gives $\alpha = \frac{A + B}{2}$ and $\beta = \frac{A - B}{2}$.

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$. This gives $\alpha = \frac{A + B}{2}$ and $\beta = \frac{A - B}{2}$. So we get the following formula:

$$\sin A + \sin B = 2 \sin\left(\frac{A + B}{2}\right) \cos\left(\frac{A - B}{2}\right)$$

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$. This gives $\alpha = \frac{A + B}{2}$ and $\beta = \frac{A - B}{2}$. So we get the following formula:

$$\sin A + \sin B = 2 \sin\left(\frac{A + B}{2}\right) \cos\left(\frac{A - B}{2}\right)$$

Similarly if we subtracted the second equation from the first one we would get:

$$\sin A - \sin B = 2 \sin\left(\frac{A - B}{2}\right) \cos\left(\frac{A + B}{2}\right)$$

We can also add the third and fourth equations to get:

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta$$

and again using the substitutions $A = \alpha + \beta$ and $B = \alpha - \beta$ we get:

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

We can also add the third and fourth equations to get:

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta$$

and again using the substitutions $A = \alpha + \beta$ and $B = \alpha - \beta$ we get:

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

Subtracting the fourth equation from the third one gives:

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

In the end we get the following four formulae:

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\sin A - \sin B = 2 \sin\left(\frac{A-B}{2}\right) \cos\left(\frac{A+B}{2}\right)$$

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

In the end we get the following four formulae:

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\sin A - \sin B = 2 \sin\left(\frac{A-B}{2}\right) \cos\left(\frac{A+B}{2}\right)$$

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

These formulae are in the formula booklet but you should remember them nevertheless.

Exercise 1

Evaluate $\sin 105^\circ - \sin 15^\circ$.

Exercise 1

Evaluate $\sin 105^\circ - \sin 15^\circ$.

$$\sin 105^\circ - \sin 15^\circ = 2 \sin 45^\circ \cos 60^\circ = 2 \times \frac{\sqrt{2}}{2} \times \frac{1}{2} = \frac{\sqrt{2}}{2}$$

Exercise 2

Evaluate $\cos 105^\circ - \sin 75^\circ$.

Exercise 2

Evaluate $\cos 105^\circ - \sin 75^\circ$. (Remember about the formula that changes a function into a co-function).

Exercise 2

Evaluate $\cos 105^\circ - \sin 75^\circ$. (Remember about the formula that changes a function into a co-function).

$$\cos 105^\circ - \sin 75^\circ = \cos 105^\circ - \cos 15^\circ = -2 \sin 60^\circ \sin 45^\circ = -\frac{\sqrt{6}}{2}$$

Przypomnienie

Teraz przejdziemy do rozwiązywania równań.

Przypomnienie

Teraz przejdziemy do rozwiązywania równań. Krótkie przypomnienie.

Przypomnienie

Teraz przejdziemy do rozwiązywania równań. Krótkie przypomnienie.
Równanie

$$\sin x = \frac{1}{2}$$

rozwiązuje my korzystając z wykresu lub okręgu jednostkowego i dostajemy:

$$x = \frac{\pi}{6} + 2k\pi \quad \vee \quad x = \frac{5\pi}{6} + 2k\pi$$

gdzie $k \in \mathbb{Z}$.

Przypomnienie

Jeśli zamiast $\sin x$ mamy np. $\sin 10x$ to rozwiązuje się analogicznie:

$$\sin 10x = \frac{1}{2}$$

teraz mamy:

$$10x = \frac{\pi}{6} + 2k\pi \quad \vee \quad 10x = \frac{5\pi}{6} + 2k\pi$$

czyli:

$$x = \frac{\pi}{60} + \frac{1}{5}k\pi \quad \vee \quad x = \frac{\pi}{12} + \frac{1}{5}k\pi$$

gdzie $k \in \mathbb{Z}$.

Rozwiązywanie równań trygonometrycznych będziemy jeszcze intensywnie ćwiczyli. Poniższe przykłady pokażą, jak możemy nasze nowe wzory do tego wykorzystać.

Exercise 3

Express $\sin \theta + \sin 3\theta$ as a product of two trigonometric functions. Hence solve $\sin \theta + \sin 2\theta + \sin 3\theta = 0$ for $0 \leq \theta \leq \pi$.

Exercise 3

Express $\sin \theta + \sin 3\theta$ as a product of two trigonometric functions. Hence solve $\sin \theta + \sin 2\theta + \sin 3\theta = 0$ for $0 \leq \theta \leq \pi$.

We have:

$$\begin{aligned}\sin \theta + \sin 3\theta &= 2 \sin\left(\frac{\theta + 3\theta}{2}\right) \cos\left(\frac{\theta - 3\theta}{2}\right) = \\ &= 2 \sin 2\theta \cos(-\theta) = 2 \sin 2\theta \cos \theta\end{aligned}$$

Exercise 3

Express $\sin \theta + \sin 3\theta$ as a product of two trigonometric functions. Hence solve $\sin \theta + \sin 2\theta + \sin 3\theta = 0$ for $0 \leq \theta \leq \pi$.

We have:

$$\begin{aligned}\sin \theta + \sin 3\theta &= 2 \sin\left(\frac{\theta + 3\theta}{2}\right) \cos\left(\frac{\theta - 3\theta}{2}\right) = \\ &= 2 \sin 2\theta \cos(-\theta) = 2 \sin 2\theta \cos \theta\end{aligned}$$

So we get:

$$\begin{aligned}\sin \theta + \sin 2\theta + \sin 3\theta &= 0 \\ 2 \sin 2\theta \cos \theta + \sin 2\theta &= 0 \\ \sin 2\theta(2 \cos \theta + 1) &= 0\end{aligned}$$

Which gives $\sin 2\theta = 0$ or $2 \cos \theta + 1 = 0$

Exercise 3

We want to solve $\sin 2\theta = 0$ or $2\cos \theta + 1 = 0$ for $0 \leq \theta \leq \pi$.

Exercise 3

We want to solve $\sin 2\theta = 0$ or $2\cos \theta + 1 = 0$ for $0 \leq \theta \leq \pi$.

The first equation gives us $\theta \in \{0, \frac{\pi}{2}, \pi\}$, the second gives us $\theta = \frac{2\pi}{3}$, so in the end we have four solutions: $0, \frac{\pi}{2}, \frac{2\pi}{3}, \pi$.

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$.

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$. We get:

$$\begin{aligned}\cos 5\theta - \cos \theta &= -2 \sin\left(\frac{5\theta + \theta}{2}\right) \sin\left(\frac{5\theta - \theta}{2}\right) = \\ &= -2 \sin 3\theta \sin 2\theta\end{aligned}$$

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$. We get:

$$\begin{aligned}\cos 5\theta - \cos \theta &= -2 \sin\left(\frac{5\theta + \theta}{2}\right) \sin\left(\frac{5\theta - \theta}{2}\right) = \\ &= -2 \sin 3\theta \sin 2\theta\end{aligned}$$

Hence we get:

$$\cos 5\theta - \cos \theta = \sin 2\theta \quad \Rightarrow \quad -2 \sin 3\theta \sin 2\theta = \sin 2\theta$$

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$. We get:

$$\begin{aligned}\cos 5\theta - \cos \theta &= -2 \sin\left(\frac{5\theta + \theta}{2}\right) \sin\left(\frac{5\theta - \theta}{2}\right) = \\ &= -2 \sin 3\theta \sin 2\theta\end{aligned}$$

Hence we get:

$$\cos 5\theta - \cos \theta = \sin 2\theta \quad \Rightarrow \quad -2 \sin 3\theta \sin 2\theta = \sin 2\theta$$

This gives:

$$\sin 2\theta + 2 \sin 3\theta \sin 2\theta = 0$$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

So

$$\sin 2\theta = 0 \quad \text{or} \quad \sin 3\theta = -\frac{1}{2}$$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

So

$$\sin 2\theta = 0 \quad \text{or} \quad \sin 3\theta = -\frac{1}{2}$$

From the first equation we get $\theta \in \{0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi\}$. The second equation gives $\theta \in \{\frac{7\pi}{18}, \frac{11\pi}{18}, \frac{19\pi}{18}, \frac{23\pi}{18}, \frac{31\pi}{18}, \frac{35\pi}{18}\}$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

So

$$\sin 2\theta = 0 \quad \text{or} \quad \sin 3\theta = -\frac{1}{2}$$

From the first equation we get $\theta \in \{0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi\}$. The second equation gives $\theta \in \{\frac{7\pi}{18}, \frac{11\pi}{18}, \frac{19\pi}{18}, \frac{23\pi}{18}, \frac{31\pi}{18}, \frac{35\pi}{18}\}$

So in the end we have 11 solutions:

$$\theta \in \{0, \frac{7\pi}{18}, \frac{\pi}{2}, \frac{11\pi}{18}, \pi, \frac{19\pi}{18}, \frac{23\pi}{18}, \frac{3\pi}{2}, \frac{31\pi}{18}, \frac{35\pi}{18}, 2\pi\}$$

In case of any questions you can email me at T.J.Lechowski@gmail.com.