Probability distribution 17.01 [52 marks]

A game is played where two unbiased dice are rolled and the score in the game is the greater of the two numbers shown. If the two numbers are the same, then the score in the game is the number shown on one of the dice. A diagram showing the possible outcomes is given below.

		I			t die		
		1	2	3	4	5	6
_	1	•	-	•	•	•	•
	2	•	•	•	•	•	•
Second die	3	•	•	•	•	•	•
Secol	4	•	•	•	•	•	٠
	5	•	•	•	•	•	٠
	6	•	•	•	•	•	•

Let T be the random variable "the score in a game".

1a. Complete the table to show the probability distribution of T.

[2 marks]

t	1	2	3	4	5	6
$\mathbf{P}(T=t)$						

Markscheme

t	1	2	3	4	5	6	
$\mathbf{P}(T=t)$	1 36 (0.027777)	3 36 (0.083333)	5 36 (0.138888)	7 36 (0.194444)	9 36 (0.25)	11 36 (0.305555)	A2

Note: Award **A1** if three to five probabilities are correct.

Find the probability that

1b. a player scores at least 3 in a game.




1c. a player scores 6, given that they scored at least 3.

Markscheme use of conditional probability (M1) e.g. denominator of 32 **OR** denominator of 0. 8888888..., etc. $\frac{11}{32}(0.34375, 34.4\%)$ A1 [2 marks]

1d. Find the expected score of a game.

[2 marks]



[1 mark]

Jae Hee plays a game involving a biased six-sided die.

The faces of the die are labelled -3, -1, 0, 1, 2 and 5.

The score for the game, X, is the number which lands face up after the die is rolled.

The following table shows the probability distribution for X.

Score x	-3	-1	0	1	2	5
P(X=x)	$\frac{1}{18}$	р	$\frac{3}{18}$	$\frac{1}{18}$	$\frac{2}{18}$	$\frac{7}{18}$

2a. Find the exact value of p.

Mar	kscheme
$\frac{4}{18}\left(\frac{2}{9}\right)$	A1
[1 mark	1

Jae Hee plays the game once.

2b. Calculate the expected score.

Markscheme
$-3 imes rac{1}{18} + (-1) imes rac{4}{18} + 0 imes rac{3}{18} + \ldots + 5 imes rac{7}{18}$ (M1)
Note: Award <i>(M1)</i> for their correct substitution into the formula for expected value.
$= 1.83 \left(rac{33}{18}, 1.83333 \ldots ight)$ Al
[2 marks]

2c. Jae Hee plays the game twice and adds the two scores together.[3 marks]Find the probability Jae Hee has a **total** score of -3.

[1 mark]

Markscheme $2 imesrac{1}{18} imesrac{3}{18}$ (M1)(M1) **Note:** Award *(M1)* for $\frac{1}{18} \times \frac{3}{18}$, award *(M1)* for multiplying their product by 2. $=rac{1}{54} igg(rac{6}{324}, 0.0185185 \ldots, 1.85\% igg)$ A1 [3 marks]

There are three fair six-sided dice. Each die has two green faces, two yellow faces and two red faces.

diagram)

All three dice are rolled.

3a. Find the probability of rolling exactly one red face.

```
[2 marks]
```

Markscheme
valid approach to find P(one red) (M1)
eg $_{n}C_{a} imes p^{a} imes q^{n-a}$, $\mathrm{B}\left(n,p ight)$, $3\left(rac{1}{3} ight)\left(rac{2}{3} ight)^{2}$, $\left(rac{3}{1} ight)$
listing all possible cases for exactly one red (may be indicated on tree
P(1 red) = 0.444 $\left(=\frac{4}{9}\right)$ [0.444, 0.445] A1 N2
[3 marks] [5 maximum for parts (a.i) and (a.ii)]

3b. Find the probability of rolling two or more red faces.

[3 marks]

valid approach (M1) eg P(X = 2) + P(X = 3), 1 - P(X \le 1), binomcdf $(3, \frac{1}{3}, 2, 3)$ correct working (A1) eg $\frac{2}{9} + \frac{1}{27}$, 0.222 + 0.037, $1 - (\frac{2}{3})^3 - \frac{4}{9}$ 0.259259 P(at least two red) = $0.259(=\frac{7}{27})$ A1 N3 [3 marks] [5 maximum for parts (a.i) and (a.ii)]

Ted plays a game using these dice. The rules are:

- Having a turn means to roll all three dice.
- He wins \$10 for each green face rolled and adds this to his winnings.
- After a turn Ted can either:
 end the game (and keep his winnings), or
 have another turn (and try to increase his winnings).
- If two or more red faces are rolled in a turn, all winnings are lost and the game ends.
- 3c. Show that, after a turn, the probability that Ted adds exactly \$10 to his [5 marks] winnings is $\frac{1}{3}$.

recognition that winning \$10 means rolling exactly one green(M1)recognition that winning \$10 also means rolling at most 1 red(M1)

eg "cannot have 2 or more reds"

correct approach **A1**

 $eg P(1G \cap 0R) + P(1G \cap 1R), P(1G) - P(1G \cap 2R),$

"one green and two yellows or one of each colour"

Note: Because this is a "show that" question, do not award this **A1** for purely numerical expressions.

one correct probability for their approach (A1)

eg $3\left(\frac{1}{3}\right)\left(\frac{1}{3}\right)^2$, $\frac{6}{27}$, $3\left(\frac{1}{3}\right)\left(\frac{2}{3}\right)^2$, $\frac{1}{9}$, $\frac{2}{9}$ correct working leading to $\frac{1}{3}$ **A1** *eg* $\frac{3}{27} + \frac{6}{27}$, $\frac{12}{27} - \frac{3}{27}$, $\frac{1}{9} + \frac{2}{9}$ probability = $\frac{1}{3}$ **AG NO** [5 marks]

The random variable D (\$) represents how much is added to his winnings after a turn.

The following table shows the distribution for D, where $\$ represents his winnings in the game so far.

D (\$)	-w	0	10	20	30
$\mathbb{P}\left(D=d\right)$	x	у	$\frac{1}{3}$	$\frac{2}{9}$	$\frac{1}{27}$

3d. Write down the value of x.

[1 mark]

Markscheme	
$x=rac{7}{27}$, 0.259 (check FT from (a)(ii))	A1 N1
[1 mark]	

Markscheme evidence of summing probabilities to 1 **(M1)** $eg \ \sum = 1, \ x + y + \frac{1}{3} + \frac{2}{9} + \frac{1}{27} = 1, \ 1 - \frac{7}{27} - \frac{9}{27} - \frac{6}{27} - \frac{1}{27}$ 0.148147 (0.148407 if working with **their** *x* value to 3 sf) $y = \frac{4}{27}$ (exact), 0.148 **A1 N2** [2 marks]

3f. Ted will always have another turn if he expects an increase to his [3 marks] winnings.

Find the least value of \boldsymbol{w} for which Ted should end the game instead of having another turn.


```
Markscheme
correct substitution into the formula for expected value (A1)
eg - w \cdot \frac{7}{27} + 10 \cdot \frac{9}{27} + 20 \cdot \frac{6}{27} + 30 \cdot \frac{1}{27}
correct critical value (accept inequality) A1
eg w = 34.2857 (= \frac{240}{7}), w > 34.2857
$40 A1 N2
[3 marks]
```

A bag contains n marbles, two of which are blue. Hayley plays a game in which she randomly draws marbles out of the bag, one after another, without replacement. The game ends when Hayley draws a blue marble.

4a. Find the probability, in terms of *n*, that the game will end on her first [1 mark] draw.



4b. Find the probability, in terms of n, that the game will end on her second [3 marks] draw.



Let n = 5. Find the probability that the game will end on her

4c. third draw.



[2 marks]

Markscheme
correct working (A1)
eg $rac{3}{5} imesrac{2}{4} imesrac{1}{3} imesrac{2}{2}$
$rac{6}{60}~\left(=rac{1}{10} ight)$ A1 N2
[2 marks]

4e. Hayley plays the game when n = 5. She pays \$20 to play and can earn [7 marks] money back depending on the number of draws it takes to obtain a blue marble. She earns no money back if she obtains a blue marble on her first draw. Let M be the amount of money that she earns back playing the game. This information is shown in the following table.

Number of draws	1	2	3	4
Money earned back (\$ <i>M</i>)	0	20	<u>8k</u>	12k

Find the value of k so that this is a fair game.

correct probabilities (seen anywhere) (A1)(A1) eg $P(1) = \frac{2}{5}$, $P(2) = \frac{6}{20}$ (may be seen on tree diagram) valid approach to find E (M) or expected winnings using their probabilities (M1) eq $P(1) \times (0) + P(2) \times (20) + P(3) \times (8k) + P(4) \times (12k)$, $P(1) \times (-20) + P(2) \times (0) + P(3) \times (8k - 20) + P(4) \times (12k - 20)$ correct working to find E (*M*) or expected winnings (A1) $eg \frac{2}{5}(0) + \frac{3}{10}(20) + \frac{1}{5}(8k) + \frac{1}{10}(12k),$ $\frac{2}{5}(-20) + \frac{3}{10}(0) + \frac{1}{5}(8k - 20) + \frac{1}{10}(12k - 20)$ correct equation for fair game A1 eg $\frac{3}{10}(20) + \frac{1}{5}(8k) + \frac{1}{10}(12k) = 20$, $\frac{2}{5}(-20) + \frac{1}{5}(8k - 20) + \frac{1}{10}(12k - 20) = 0$ correct working to combine terms in k(A1) eg $-8 + \frac{14}{5}k - 4 - 2 = 0$, $6 + \frac{14}{5}k = 20$, $\frac{14}{5}k = 14$ *k* = 5 *A1 N0* Note: Do not award the final A1 if the candidate's FT probabilities do not sum to 1.

[7 marks]

A discrete random variable X has the following probability distribution.

Х	0	1	2	3
P(X=x)	0.475	$2k^2$	$\frac{k}{10}$	6 <i>k</i> ²

5a. Find the value of k.

[4 marks]

* This question is from an exam for a previous syllabus, and may contain minor differences in marking or structure.

valid approach **(M1)** *eg*total probability = 1 correct equation **(A1)** *eg* $0.475 + 2k^2 + \frac{k}{10} + 6k^2 = 1, 8k^2 + 0.1k - 0.525 = 0$ k = 0.25 **A2 N3 [4 marks]**

5b. Write down P(X = 2).

 Markscheme

 P(X = 2) = 0.025 A1
 N1

 [1 mark]
 [1 mark]

[1 mark]

[3 marks]

5c. Find
$$P(X = 2|X > 0)$$
.

Markscheme valid approach for finding P(X > 0) (M1) $eg1 - 0.475, 2(0.25^2) + 0.025 + 6(0.25^2), 1 - P(X = 0), 2k^2 + \frac{k}{10} + 6k^2$ correct substitution into formula for conditional probability (A1) $eg\frac{0.025}{1-0.475}, \frac{0.025}{0.525}$ 0.0476190 $P(X = 2|X > 0) = \frac{1}{21}$ (exact), 0.0476 A1 N2 [3 marks]



Printed for 2 SPOLECZNE LICEUM