

Sum/difference - to - product formulae

When we solve an equation like:

$$x^2 - 4x - 12 = 0 \quad (1)$$

what we want to do is to turn it (if possible) into:

$$(x - 6)(x + 2) = 0 \quad (2)$$

because this immediately gives us the solutions $x = 6$ or $x = -2$.

When we solve an equation like:

$$x^2 - 4x - 12 = 0 \quad (1)$$

what we want to do is to turn it (if possible) into:

$$(x - 6)(x + 2) = 0 \quad (2)$$

because this immediately gives us the solutions $x = 6$ or $x = -2$.

We turned equation (1) where we add terms to an equation (2) where we multiply terms. This is often useful when solving equations.

We want to do something similar to trigonometric functions. Suppose we have an equation:

$$\sin A + \cos B = 0$$

can we somehow turn this sum into a product?

We want to do something similar to trigonometric functions. Suppose we have an equation:

$$\sin A + \cos B = 0$$

can we somehow turn this sum into a product?

The answer is of course yes and we will see how to do this on the next few slides.

Recall that we already have the following four formulae:

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \sin \beta \cos \alpha$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$.

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$. This gives $\alpha = \frac{A + B}{2}$ and $\beta = \frac{A - B}{2}$.

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$. This gives $\alpha = \frac{A + B}{2}$ and $\beta = \frac{A - B}{2}$. So we get the following formula:

$$\sin A + \sin B = 2 \sin\left(\frac{A + B}{2}\right) \cos\left(\frac{A - B}{2}\right)$$

If we add the first two equations we get:

$$\sin(\alpha + \beta) + \sin(\alpha - \beta) = 2 \sin \alpha \cos \beta$$

Now let $A = \alpha + \beta$ and $B = \alpha - \beta$. This gives $\alpha = \frac{A + B}{2}$ and $\beta = \frac{A - B}{2}$. So we get the following formula:

$$\sin A + \sin B = 2 \sin\left(\frac{A + B}{2}\right) \cos\left(\frac{A - B}{2}\right)$$

Similarly if we subtracted the second equation from the first one we would get:

$$\sin A - \sin B = 2 \sin\left(\frac{A - B}{2}\right) \cos\left(\frac{A + B}{2}\right)$$

We can also add the third and fourth equations to get:

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta$$

and again using the substitutions $A = \alpha + \beta$ and $B = \alpha - \beta$ we get:

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

We can also add the third and fourth equations to get:

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta$$

and again using the substitutions $A = \alpha + \beta$ and $B = \alpha - \beta$ we get:

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

Subtracting the fourth equation from the third one gives:

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

In the end we get the following four formulae:

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\sin A - \sin B = 2 \sin\left(\frac{A-B}{2}\right) \cos\left(\frac{A+B}{2}\right)$$

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

In the end we get the following four formulae:

$$\sin A + \sin B = 2 \sin\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\sin A - \sin B = 2 \sin\left(\frac{A-B}{2}\right) \cos\left(\frac{A+B}{2}\right)$$

$$\cos A + \cos B = 2 \cos\left(\frac{A+B}{2}\right) \cos\left(\frac{A-B}{2}\right)$$

$$\cos A - \cos B = -2 \sin\left(\frac{A+B}{2}\right) \sin\left(\frac{A-B}{2}\right)$$

The IB does not require you to know these, but they are very helpful. I would highly recommend to learn them by heart (and since they're not required by the IB, they're not in the formula booklet).

Exercise 1

Evaluate $\sin 105^\circ - \sin 15^\circ$.

Exercise 1

Evaluate $\sin 105^\circ - \sin 15^\circ$.

$$\sin 105^\circ - \sin 15^\circ = 2 \sin 45^\circ \cos 60^\circ = 2 \times \frac{\sqrt{2}}{2} \times \frac{1}{2} = \frac{\sqrt{2}}{2}$$

Exercise 2

Evaluate $\cos 105^\circ - \sin 75^\circ$.

Exercise 2

Evaluate $\cos 105^\circ - \sin 75^\circ$.

We first need to change \sin into \cos . We can do so using the reduction formulae $\sin(90^\circ - \alpha) = \cos \alpha$:

Exercise 2

Evaluate $\cos 105^\circ - \sin 75^\circ$.

We first need to change \sin into \cos . We can do so using the reduction formulae $\sin(90^\circ - \alpha) = \cos \alpha$:

$$\cos 105^\circ - \sin 75^\circ = \cos 105^\circ - \cos 15^\circ = -2 \sin 60^\circ \sin 45^\circ = -\frac{\sqrt{6}}{2}$$

Exercise 3

Express $\sin \theta + \sin 3\theta$ as a product of two trigonometric functions. Hence solve $\sin \theta + \sin 2\theta + \sin 3\theta = 0$ for $0 \leq \theta \leq \pi$.

Exercise 3

Express $\sin \theta + \sin 3\theta$ as a product of two trigonometric functions. Hence solve $\sin \theta + \sin 2\theta + \sin 3\theta = 0$ for $0 \leq \theta \leq \pi$.

We have:

$$\begin{aligned}\sin \theta + \sin 3\theta &= 2 \sin\left(\frac{\theta + 3\theta}{2}\right) \cos\left(\frac{\theta - 3\theta}{2}\right) = \\ &= 2 \sin 2\theta \cos(-\theta) = 2 \sin 2\theta \cos \theta\end{aligned}$$

Exercise 3

Express $\sin \theta + \sin 3\theta$ as a product of two trigonometric functions. Hence solve $\sin \theta + \sin 2\theta + \sin 3\theta = 0$ for $0 \leq \theta \leq \pi$.

We have:

$$\begin{aligned}\sin \theta + \sin 3\theta &= 2 \sin\left(\frac{\theta + 3\theta}{2}\right) \cos\left(\frac{\theta - 3\theta}{2}\right) = \\ &= 2 \sin 2\theta \cos(-\theta) = 2 \sin 2\theta \cos \theta\end{aligned}$$

So we get:

$$\begin{aligned}\sin \theta + \sin 2\theta + \sin 3\theta &= 0 \\ 2 \sin 2\theta \cos \theta + \sin 2\theta &= 0 \\ \sin 2\theta(2 \cos \theta + 1) &= 0\end{aligned}$$

Which gives $\sin 2\theta = 0$ or $2 \cos \theta + 1 = 0$

Exercise 3

We want to solve $\sin 2\theta = 0$ or $2 \cos \theta + 1 = 0$ for $0 \leq \theta \leq \pi$.

Exercise 3

We want to solve $\sin 2\theta = 0$ or $2 \cos \theta + 1 = 0$ for $0 \leq \theta \leq \pi$.

The first equation gives us $\theta \in \{0, \frac{\pi}{2}, \pi\}$, the second gives us $\theta = \frac{2\pi}{3}$, so in the end we have four solutions: $0, \frac{\pi}{2}, \frac{2\pi}{3}, \pi$.

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$.

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$. We get:

$$\begin{aligned}\cos 5\theta - \cos \theta &= -2 \sin\left(\frac{5\theta + \theta}{2}\right) \sin\left(\frac{5\theta - \theta}{2}\right) = \\ &= -2 \sin 3\theta \sin 2\theta\end{aligned}$$

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$. We get:

$$\begin{aligned}\cos 5\theta - \cos \theta &= -2 \sin\left(\frac{5\theta + \theta}{2}\right) \sin\left(\frac{5\theta - \theta}{2}\right) = \\ &= -2 \sin 3\theta \sin 2\theta\end{aligned}$$

Hence we get:

$$\cos 5\theta - \cos \theta = \sin 2\theta \quad \Rightarrow \quad -2 \sin 3\theta \sin 2\theta = \sin 2\theta$$

Exercise 4

Solve $\cos 5\theta - \cos \theta = \sin 2\theta$ for $0 \leq \theta \leq 2\pi$.

We start by combining $\cos 5\theta - \cos \theta$. We get:

$$\begin{aligned}\cos 5\theta - \cos \theta &= -2 \sin\left(\frac{5\theta + \theta}{2}\right) \sin\left(\frac{5\theta - \theta}{2}\right) = \\ &= -2 \sin 3\theta \sin 2\theta\end{aligned}$$

Hence we get:

$$\cos 5\theta - \cos \theta = \sin 2\theta \quad \Rightarrow \quad -2 \sin 3\theta \sin 2\theta = \sin 2\theta$$

This gives:

$$\sin 2\theta + 2 \sin 3\theta \sin 2\theta = 0$$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

So

$$\sin 2\theta = 0 \quad \text{or} \quad \sin 3\theta = -\frac{1}{2}$$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

So

$$\sin 2\theta = 0 \quad \text{or} \quad \sin 3\theta = -\frac{1}{2}$$

From the first equation we get $\theta \in \{0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi\}$. The second equation gives $\theta \in \{\frac{7\pi}{18}, \frac{11\pi}{18}, \frac{19\pi}{18}, \frac{23\pi}{18}, \frac{31\pi}{18}, \frac{35\pi}{18}\}$

Exercise 4

Factoring $\sin 2\theta$ gives:

$$\sin 2\theta(1 + 2 \sin 3\theta) = 0$$

So

$$\sin 2\theta = 0 \quad \text{or} \quad \sin 3\theta = -\frac{1}{2}$$

From the first equation we get $\theta \in \{0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi\}$. The second equation gives $\theta \in \{\frac{7\pi}{18}, \frac{11\pi}{18}, \frac{19\pi}{18}, \frac{23\pi}{18}, \frac{31\pi}{18}, \frac{35\pi}{18}\}$

So in the end we have 11 solutions:

$$\theta \in \{0, \frac{7\pi}{18}, \frac{\pi}{2}, \frac{11\pi}{18}, \pi, \frac{19\pi}{18}, \frac{23\pi}{18}, \frac{3\pi}{2}, \frac{31\pi}{18}, \frac{35\pi}{18}, 2\pi\}$$

In case of any questions you can email me at T.J.Lechowski@gmail.com.