

Disguised quadratic equations

Introduction

Before you start make sure you are comfortable with solving quadratic equations using factorization, completing the square or quadratic formula and that you are able recognize when a quadratic equation has no solutions.

Introduction

Sometimes a complicated equation can be reduced to a quadratic equation by introducing a new variable.

Introduction

Sometimes a complicated equation can be reduced to a quadratic equation by introducing a new variable. Consider for instance:

$$\left(\frac{1}{x+1}\right)^2 - \frac{3}{x+1} - 10 = 0$$

Introduction

Sometimes a complicated equation can be reduced to a quadratic equation by introducing a new variable. Consider for instance:

$$\left(\frac{1}{x+1}\right)^2 - \frac{3}{x+1} - 10 = 0$$

This may look like a complicated equation, but in fact it can be easily reduced to a quadratic, which we can solve in few seconds.

Introduction

$$\left(\frac{1}{x+1}\right)^2 - \frac{3}{x+1} - 10 = 0$$

We need to start with the assumption that $x \neq -1$, because we don't want to have 0 in the denominator.

Introduction

$$\left(\frac{1}{x+1}\right)^2 - \frac{3}{x+1} - 10 = 0$$

We need to start with the assumption that $x \neq -1$, because we don't want to have 0 in the denominator. Now we can introduce an auxiliary variable.

We will let $t = \frac{1}{x+1}$. If we now substitute t into our equation, we get:

Introduction

$$\left(\frac{1}{x+1}\right)^2 - \frac{3}{x+1} - 10 = 0$$

We need to start with the assumption that $x \neq -1$, because we don't want to have 0 in the denominator. Now we can introduce an auxiliary variable.

We will let $t = \frac{1}{x+1}$. If we now substitute t into our equation, we get:

$$t^2 - 3t - 10 = 0$$

Introduction

$$\left(\frac{1}{x+1}\right)^2 - \frac{3}{x+1} - 10 = 0$$

We need to start with the assumption that $x \neq -1$, because we don't want to have 0 in the denominator. Now we can introduce an auxiliary variable.

We will let $t = \frac{1}{x+1}$. If we now substitute t into our equation, we get:

$$t^2 - 3t - 10 = 0$$

This can be easily solved using factorization:

$$(t - 5)(t + 2) = 0$$

$$t - 5 = 0 \quad \text{or} \quad t + 2 = 0$$

$$t = 5 \quad \text{or} \quad t = -2$$

Introduction

Remember however that no one asked us about t !

Introduction

Remember however that no one asked us about t ! We need to go back to our substitution and solve for x . Remember $t = \frac{1}{x+1}$, so we have:

Introduction

Remember however that no one asked us about t ! We need to go back to our substitution and solve for x . Remember $t = \frac{1}{x+1}$, so we have:

$$\begin{array}{lcl} \frac{1}{x+1} = 5 & \text{or} & \frac{1}{x+1} = -2 \\ 1 = 5x + 5 & \text{or} & 1 = -2x - 2 \\ x = -\frac{4}{5} & \text{or} & x = -\frac{3}{2} \end{array}$$

Introduction

Remember however that no one asked us about t ! We need to go back to our substitution and solve for x . Remember $t = \frac{1}{x+1}$, so we have:

$$\begin{array}{lcl} \frac{1}{x+1} = 5 & \text{or} & \frac{1}{x+1} = -2 \\ 1 = 5x + 5 & \text{or} & 1 = -2x - 2 \\ x = -\frac{4}{5} & \text{or} & x = -\frac{3}{2} \end{array}$$

And these are our final two solutions.

What we need to practice now is the ability to recognize when a seemingly complicated equation can be reduced to a quadratic by introducing a new variable.

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Here setting $t = x^3$ reduces the above equation to a simple quadratic:

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Here setting $t = x^3$ reduces the above equation to a simple quadratic:

$$t^2 - 10t + 16 = 0$$

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Here setting $t = x^3$ reduces the above equation to a simple quadratic:

$$t^2 - 10t + 16 = 0$$

We can now factorize the left hand side and solve:

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Here setting $t = x^3$ reduces the above equation to a simple quadratic:

$$t^2 - 10t + 16 = 0$$

We can now factorize the left hand side and solve:

$$(t - 8)(t - 2) = 0$$

$$t - 8 = 0 \quad \text{or} \quad t - 2 = 0$$

$$t = 8 \quad \text{or} \quad t = 2$$

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Here setting $t = x^3$ reduces the above equation to a simple quadratic:

$$t^2 - 10t + 16 = 0$$

We can now factorize the left hand side and solve:

$$(t - 8)(t - 2) = 0$$

$$t - 8 = 0 \quad \text{or} \quad t - 2 = 0$$

$$t = 8 \quad \text{or} \quad t = 2$$

Going back to x we get:

Example 1

Solve:

$$x^6 - 10x^3 + 16 = 0$$

Here setting $t = x^3$ reduces the above equation to a simple quadratic:

$$t^2 - 10t + 16 = 0$$

We can now factorize the left hand side and solve:

$$(t - 8)(t - 2) = 0$$

$$t - 8 = 0 \quad \text{or} \quad t - 2 = 0$$

$$t = 8 \quad \text{or} \quad t = 2$$

Going back to x we get:

$$x^3 = 8 \quad \text{or} \quad x^3 = 2$$

$$x = 2 \quad \text{or} \quad x = \sqrt[3]{2}$$

And these are our solutions to the original equation.

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

We start by making the assumption $x \geq 0$.

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

We start by making the assumption $x \geq 0$. This is because x appears under the $\sqrt{\quad}$ sign, and we cannot take square root of negative numbers (at least for now).

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

We start by making the assumption $x \geq 0$. This is because x appears under the $\sqrt{\quad}$ sign, and we cannot take square root of negative numbers (at least for now). Now we can introduce a variable $t = \sqrt{x}$ and substitute it into the equation to get:

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

We start by making the assumption $x \geq 0$. This is because x appears under the $\sqrt{\quad}$ sign, and we cannot take square root of negative numbers (at least for now). Now we can introduce a variable $t = \sqrt{x}$ and substitute it into the equation to get:

$$t^2 - t - 6 = 0$$

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

We start by making the assumption $x \geq 0$. This is because x appears under the $\sqrt{\quad}$ sign, and we cannot take square root of negative numbers (at least for now). Now we can introduce a variable $t = \sqrt{x}$ and substitute it into the equation to get:

$$t^2 - t - 6 = 0$$

This can be easily solved using factorization:

Example 2

Solve:

$$x - \sqrt{x} - 6 = 0$$

We start by making the assumption $x \geq 0$. This is because x appears under the $\sqrt{\quad}$ sign, and we cannot take square root of negative numbers (at least for now). Now we can introduce a variable $t = \sqrt{x}$ and substitute it into the equation to get:

$$t^2 - t - 6 = 0$$

This can be easily solved using factorization:

$$(t - 3)(t + 2) = 0$$

$$t - 3 = 0 \quad \text{or} \quad t + 2 = 0$$

$$t = 3 \quad \text{or} \quad t = -2$$

Example 2

Going back to x we get:

Example 2

Going back to x we get:

$$\begin{array}{lcl} \sqrt{x} = 3 & \text{or} & \sqrt{x} = -2 \\ x = 9 & \text{or} & \text{no solution} \end{array}$$

So in the end we only have one solution $x = 9$.

Example 3

Solve:

$$2(x^2 + 1)^2 - 5(x^2 + 1) - 3 = 0$$

Example 3

Solve:

$$2(x^2 + 1)^2 - 5(x^2 + 1) - 3 = 0$$

We will set $t = x^2 + 1$ and now our equation becomes:

Example 3

Solve:

$$2(x^2 + 1)^2 - 5(x^2 + 1) - 3 = 0$$

We will set $t = x^2 + 1$ and now our equation becomes:

$$2t^2 - 5t - 3 = 0$$

Example 3

Solve:

$$2(x^2 + 1)^2 - 5(x^2 + 1) - 3 = 0$$

We will set $t = x^2 + 1$ and now our equation becomes:

$$2t^2 - 5t - 3 = 0$$

We factorize and get:

Example 3

Solve:

$$2(x^2 + 1)^2 - 5(x^2 + 1) - 3 = 0$$

We will set $t = x^2 + 1$ and now our equation becomes:

$$2t^2 - 5t - 3 = 0$$

We factorize and get:

$$(2t + 1)(t - 3) = 0$$

$$2t + 1 = 0 \quad \text{or} \quad t - 3 = 0$$

$$t = -\frac{1}{2} \quad \text{or} \quad t = 3$$

Example 3

Now we go back to x we get:

Example 3

Now we go back to x we get:

$$x^2 + 1 = -\frac{1}{2} \quad \text{or} \quad x^2 + 1 = 3$$

$$x^2 = -\frac{3}{2} \quad \text{or} \quad x^2 = 2$$

$$\text{no real solutions} \quad \text{or} \quad x = \pm\sqrt{2}$$

So we have two real solution $x = \sqrt{2}$ or $x = -\sqrt{2}$.

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

We need to start with the assumption $x \geq 0$.

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

We need to start with the assumption $x \geq 0$. Now we set $t = \sqrt{x}$ and get:

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

We need to start with the assumption $x \geq 0$. Now we set $t = \sqrt{x}$ and get:

$$t^2 - 2t - 4 = 0$$

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

We need to start with the assumption $x \geq 0$. Now we set $t = \sqrt{x}$ and get:

$$t^2 - 2t - 4 = 0$$

This cannot be factorized easily so we will use the quadratic formula to solve the above:

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

We need to start with the assumption $x \geq 0$. Now we set $t = \sqrt{x}$ and get:

$$t^2 - 2t - 4 = 0$$

This cannot be factorized easily so we will use the quadratic formula to solve the above:

$$\Delta = b^2 - 4ac$$

$$\Delta = (-2)^2 - 4(1)(-4) = 20$$

Example 4

Solve:

$$x - 2\sqrt{x} - 4 = 0$$

We need to start with the assumption $x \geq 0$. Now we set $t = \sqrt{x}$ and get:

$$t^2 - 2t - 4 = 0$$

This cannot be factorized easily so we will use the quadratic formula to solve the above:

$$\Delta = b^2 - 4ac$$

$$\Delta = (-2)^2 - 4(1)(-4) = 20$$

Now we have:

$$t = \frac{-b \pm \sqrt{\Delta}}{2a}$$

$$t = \frac{2 \pm \sqrt{20}}{2} = 1 \pm \sqrt{5}$$

Example 4

We go back to x we get:

Example 4

We go back to x we get:

$$\begin{array}{ll} \sqrt{x} = 1 + \sqrt{5} & \text{or} \quad \sqrt{x} = 1 - \sqrt{5} \\ x = (1 + \sqrt{5})^2 & \text{or} \quad \text{no solution} \\ x = 6 + 2\sqrt{5} & \end{array}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Example 4

We go back to x we get:

$$\begin{array}{ll} \sqrt{x} = 1 + \sqrt{5} & \text{or} \quad \sqrt{x} = 1 - \sqrt{5} \\ x = (1 + \sqrt{5})^2 & \text{or} \quad \text{no solution} \\ x = 6 + 2\sqrt{5} & \end{array}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Important point here.

Example 4

We go back to x we get:

$$\begin{array}{ll} \sqrt{x} = 1 + \sqrt{5} & \text{or} \quad \sqrt{x} = 1 - \sqrt{5} \\ x = (1 + \sqrt{5})^2 & \text{or} \quad \text{no solution} \\ x = 6 + 2\sqrt{5} & \end{array}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Important point here. You may wonder why $\sqrt{x} = 1 - \sqrt{5}$ results in no solution.

Example 4

We go back to x we get:

$$\begin{array}{ll} \sqrt{x} = 1 + \sqrt{5} & \text{or} \quad \sqrt{x} = 1 - \sqrt{5} \\ x = (1 + \sqrt{5})^2 & \text{or} \quad \text{no solution} \\ x = 6 + 2\sqrt{5} & \end{array}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Important point here. You may wonder why $\sqrt{x} = 1 - \sqrt{5}$ results in no solution. Why didn't we just square both side?

Example 4

We go back to x we get:

$$\begin{array}{ll} \sqrt{x} = 1 + \sqrt{5} & \text{or} \quad \sqrt{x} = 1 - \sqrt{5} \\ x = (1 + \sqrt{5})^2 & \text{or} \quad \text{no solution} \\ x = 6 + 2\sqrt{5} & \end{array}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Important point here. You may wonder why $\sqrt{x} = 1 - \sqrt{5}$ results in no solution. Why didn't we just square both side? It is because $1 - \sqrt{5}$ is a negative number (< 0). So there is no number whose square root is $1 - \sqrt{5}$.

Example 4

We go back to x we get:

$$\begin{aligned} \sqrt{x} &= 1 + \sqrt{5} & \text{or} & & \sqrt{x} &= 1 - \sqrt{5} \\ x &= (1 + \sqrt{5})^2 & \text{or} & & & \text{no solution} \\ x &= 6 + 2\sqrt{5} & & & & \end{aligned}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Important point here. You may wonder why $\sqrt{x} = 1 - \sqrt{5}$ results in no solution. Why didn't we just square both side? It is because $1 - \sqrt{5}$ is a negative number (< 0). So there is no number whose square root is $1 - \sqrt{5}$. It is similar to one of the previous examples where we had $\sqrt{x} = -2$.

Example 4

We go back to x we get:

$$\begin{aligned}\sqrt{x} &= 1 + \sqrt{5} & \text{or} & & \sqrt{x} &= 1 - \sqrt{5} \\ x &= (1 + \sqrt{5})^2 & \text{or} & & & \text{no solution} \\ x &= 6 + 2\sqrt{5}\end{aligned}$$

We have only one solution $x = 6 + 2\sqrt{5}$.

Important point here. You may wonder why $\sqrt{x} = 1 - \sqrt{5}$ results in no solution. Why didn't we just square both side? It is because $1 - \sqrt{5}$ is a negative number (< 0). So there is no number whose square root is $1 - \sqrt{5}$. It is similar to one of the previous examples where we had $\sqrt{x} = -2$. There isn't a number whose square root is a negative number.

Remember to always think of what you're doing.

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

In the first equation we're looking for a number whose square root is -3 .

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

In the first equation we're looking for a number whose square root is -3 . Clearly there is no such number.

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

In the first equation we're looking for a number whose square root is -3 . Clearly there is no such number. Note that $(-3)^2 = 9$, but $\sqrt{9} \neq -3$, we have $\sqrt{9} = 3$.

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

In the first equation we're looking for a number whose square root is -3 . Clearly there is no such number. Note that $(-3)^2 = 9$, but $\sqrt{9} \neq -3$, we have $\sqrt{9} = 3$.

In the second equation we're looking for a number whose cube root is -3 .

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

In the first equation we're looking for a number whose square root is -3 . Clearly there is no such number. Note that $(-3)^2 = 9$, but $\sqrt{9} \neq -3$, we have $\sqrt{9} = 3$.

In the second equation we're looking for a number whose cube root is -3 . Now we know that $(-3)^3 = -27$ and we have $\sqrt[3]{-27} = -3$.

Remember to always think of what you're doing.

Consider the two equations:

$$\sqrt{x} = -3 \qquad \sqrt[3]{x} = -3$$

Do these equations have solutions?

In the first equation we're looking for a number whose square root is -3 . Clearly there is no such number. Note that $(-3)^2 = 9$, but $\sqrt{9} \neq -3$, we have $\sqrt{9} = 3$.

In the second equation we're looking for a number whose cube root is -3 . Now we know that $(-3)^3 = -27$ and we have $\sqrt[3]{-27} = -3$. So the equation has a solution and it's $x = -27$.

Now we go back to disguised quadratics. We will now try different examples.

Example 5

Solve:

$$(x^2 + 2x)^2 + 2(x^2 + 2x) - 15 = 0$$

Example 5

Solve:

$$(x^2 + 2x)^2 + 2(x^2 + 2x) - 15 = 0$$

we substitute $t = x^2 + 2x$ and get:

Example 5

Solve:

$$(x^2 + 2x)^2 + 2(x^2 + 2x) - 15 = 0$$

we substitute $t = x^2 + 2x$ and get:

$$t^2 + 2t - 15 = 0$$

Example 5

Solve:

$$(x^2 + 2x)^2 + 2(x^2 + 2x) - 15 = 0$$

we substitute $t = x^2 + 2x$ and get:

$$t^2 + 2t - 15 = 0$$

We factorize and get:

Example 5

Solve:

$$(x^2 + 2x)^2 + 2(x^2 + 2x) - 15 = 0$$

we substitute $t = x^2 + 2x$ and get:

$$t^2 + 2t - 15 = 0$$

We factorize and get:

$$(t - 3)(t + 5) = 0$$

$$t - 3 = 0 \quad \text{or} \quad t + 5 = 0$$

$$t = 3 \quad \text{or} \quad t = -5$$

Example 5

We go back to x we get:

Example 5

We go back to x we get:

$$\begin{array}{lcl} x^2 + 2x = 3 & \text{or} & x^2 + 2x = -5 \\ x^2 + 2x - 3 = 0 & \text{or} & x^2 + 2x + 5 = 0 \end{array}$$

Example 5

We go back to x we get:

$$\begin{array}{lcl} x^2 + 2x = 3 & \text{or} & x^2 + 2x = -5 \\ x^2 + 2x - 3 = 0 & \text{or} & x^2 + 2x + 5 = 0 \end{array}$$

We will factorize the first equation:

Example 5

We go back to x we get:

$$\begin{array}{lcl} x^2 + 2x = 3 & \text{or} & x^2 + 2x = -5 \\ x^2 + 2x - 3 = 0 & \text{or} & x^2 + 2x + 5 = 0 \end{array}$$

We will factorize the first equation:

$$\begin{array}{lcl} (x + 3)(x - 1) = 0 \\ x + 3 = 0 & \text{or} & x - 1 = 0 \\ x = -3 & \text{or} & x = 1 \end{array}$$

Example 5

We go back to x we get:

$$\begin{array}{lcl} x^2 + 2x = 3 & \text{or} & x^2 + 2x = -5 \\ x^2 + 2x - 3 = 0 & \text{or} & x^2 + 2x + 5 = 0 \end{array}$$

We will factorize the first equation:

$$\begin{array}{lcl} (x + 3)(x - 1) = 0 \\ x + 3 = 0 & \text{or} & x - 1 = 0 \\ x = -3 & \text{or} & x = 1 \end{array}$$

Now we will try with the quadratic formula for the second equation:

Example 5

We go back to x we get:

$$\begin{array}{lcl} x^2 + 2x = 3 & \text{or} & x^2 + 2x = -5 \\ x^2 + 2x - 3 = 0 & \text{or} & x^2 + 2x + 5 = 0 \end{array}$$

We will factorize the first equation:

$$\begin{array}{lcl} (x + 3)(x - 1) = 0 \\ x + 3 = 0 & \text{or} & x - 1 = 0 \\ x = -3 & \text{or} & x = 1 \end{array}$$

Now we will try with the quadratic formula for the second equation:

$$\begin{array}{l} \Delta = b^2 - 4ac \\ \Delta = (2)^2 - 4(1)(5) = -16 \end{array}$$

Example 5

We go back to x we get:

$$\begin{aligned}x^2 + 2x &= 3 & \text{or} & & x^2 + 2x &= -5 \\x^2 + 2x - 3 &= 0 & \text{or} & & x^2 + 2x + 5 &= 0\end{aligned}$$

We will factorize the first equation:

$$\begin{aligned}(x + 3)(x - 1) &= 0 \\x + 3 &= 0 & \text{or} & & x - 1 &= 0 \\x &= -3 & \text{or} & & x &= 1\end{aligned}$$

Now we will try with the quadratic formula for the second equation:

$$\begin{aligned}\Delta &= b^2 - 4ac \\ \Delta &= (2)^2 - 4(1)(5) = -16\end{aligned}$$

$\Delta < 0$, so the second equation has no real solutions.

Example 5

We go back to x we get:

$$\begin{aligned}x^2 + 2x &= 3 & \text{or} & & x^2 + 2x &= -5 \\x^2 + 2x - 3 &= 0 & \text{or} & & x^2 + 2x + 5 &= 0\end{aligned}$$

We will factorize the first equation:

$$\begin{aligned}(x + 3)(x - 1) &= 0 \\x + 3 &= 0 & \text{or} & & x - 1 &= 0 \\x &= -3 & \text{or} & & x &= 1\end{aligned}$$

Now we will try with the quadratic formula for the second equation:

$$\begin{aligned}\Delta &= b^2 - 4ac \\ \Delta &= (2)^2 - 4(1)(5) = -16\end{aligned}$$

$\Delta < 0$, so the second equation has no real solutions.

In the end we have two real solutions: $x = -3$ or $x = 1$.

Example 6

Solve:

$$2\left(x + \frac{1}{x}\right)^2 - \left(x + \frac{1}{x}\right) - 10 = 0$$

Example 6

Solve:

$$2\left(x + \frac{1}{x}\right)^2 - \left(x + \frac{1}{x}\right) - 10 = 0$$

we start with the assumption $x \neq 0$ (since we have x in the denominator).

Now we substitute $t = x + \frac{1}{x}$ and get:

Example 6

Solve:

$$2\left(x + \frac{1}{x}\right)^2 - \left(x + \frac{1}{x}\right) - 10 = 0$$

we start with the assumption $x \neq 0$ (since we have x in the denominator).

Now we substitute $t = x + \frac{1}{x}$ and get:

$$2t^2 - t - 10 = 0$$

Example 6

Solve:

$$2\left(x + \frac{1}{x}\right)^2 - \left(x + \frac{1}{x}\right) - 10 = 0$$

we start with the assumption $x \neq 0$ (since we have x in the denominator).

Now we substitute $t = x + \frac{1}{x}$ and get:

$$2t^2 - t - 10 = 0$$

We factorize and get:

Example 6

Solve:

$$2\left(x + \frac{1}{x}\right)^2 - \left(x + \frac{1}{x}\right) - 10 = 0$$

we start with the assumption $x \neq 0$ (since we have x in the denominator).

Now we substitute $t = x + \frac{1}{x}$ and get:

$$2t^2 - t - 10 = 0$$

We factorize and get:

$$(2t - 5)(t + 2) = 0$$

$$2t - 5 = 0 \quad \text{or} \quad t + 2 = 0$$

$$t = \frac{5}{2} \quad \text{or} \quad t = -2$$

Example 6

We go back to x we get:

Example 6

We go back to x we get:

$$x + \frac{1}{x} = \frac{5}{2} \quad \text{or} \quad x + \frac{1}{x} = -2$$

Example 6

We go back to x we get:

$$x + \frac{1}{x} = \frac{5}{2} \quad \text{or} \quad x + \frac{1}{x} = -2$$

Since we assumed that $x \neq 0$ we can multiply both sides of both equations $2x$ and x respectively and get:

$$\begin{aligned} 2x^2 + 2 &= 5x & \text{or} & & x^2 + 1 &= -2x \\ 2x^2 - 5x + 2 &= 0 & \text{or} & & x^2 + 2x + 1 &= 0 \end{aligned}$$

Example 6

We go back to x we get:

$$x + \frac{1}{x} = \frac{5}{2} \quad \text{or} \quad x + \frac{1}{x} = -2$$

Since we assumed that $x \neq 0$ we can multiply both sides of both equations $2x$ and x respectively and get:

$$\begin{aligned} 2x^2 + 2 &= 5x & \text{or} & & x^2 + 1 &= -2x \\ 2x^2 - 5x + 2 &= 0 & \text{or} & & x^2 + 2x + 1 &= 0 \end{aligned}$$

Each equation can be solve by factorization:

Example 6

We go back to x we get:

$$x + \frac{1}{x} = \frac{5}{2} \quad \text{or} \quad x + \frac{1}{x} = -2$$

Since we assumed that $x \neq 0$ we can multiply both sides of both equations $2x$ and x respectively and get:

$$\begin{aligned} 2x^2 + 2 &= 5x & \text{or} & & x^2 + 1 &= -2x \\ 2x^2 - 5x + 2 &= 0 & \text{or} & & x^2 + 2x + 1 &= 0 \end{aligned}$$

Each equation can be solve by factorization:

$$\begin{aligned} (2x - 1)(x - 2) &= 0 & \text{or} & & (x + 1)^2 &= 0 \\ x = \frac{1}{2} & \text{or} & x = 2 & \text{or} & x = -1 \end{aligned}$$

Example 6

We go back to x we get:

$$x + \frac{1}{x} = \frac{5}{2} \quad \text{or} \quad x + \frac{1}{x} = -2$$

Since we assumed that $x \neq 0$ we can multiply both sides of both equations $2x$ and x respectively and get:

$$\begin{aligned} 2x^2 + 2 &= 5x & \text{or} & & x^2 + 1 &= -2x \\ 2x^2 - 5x + 2 &= 0 & \text{or} & & x^2 + 2x + 1 &= 0 \end{aligned}$$

Each equation can be solve by factorization:

$$\begin{aligned} (2x - 1)(x - 2) &= 0 & \text{or} & & (x + 1)^2 &= 0 \\ x = \frac{1}{2} & \text{or} & x = 2 & \text{or} & x = -1 \end{aligned}$$

We end up with three solution $x = \frac{1}{2}$ or $x = 2$ or $x = -1$.

If you have any questions you can contact me at t.j.lechowski@gmail.com.