

# Markscheme

November 2024

**Mathematics: applications and  
interpretation**

**Higher level**

**Paper 2**

© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

## Instructions to Examiners

### Abbreviations

- M** Marks awarded for attempting to use a correct **Method**.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- R** Marks awarded for clear **Reasoning**.
- AG** Answer given in the question and so no marks are awarded.
- FT** Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

### Using the markscheme

#### 1 General

Award marks using the annotations as noted in the markscheme eg **M1**, **A2**.

#### 2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies **A3**, **M2** etc., do **not** split the marks, unless there is a note.
- The response to a “show that” question does not need to restate the **AG** line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award **FT** marks as appropriate but do not award the final **A1** in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	$8\sqrt{2}$	5.65685... (incorrect decimal value)	No. Last part in question.	Award <b>A1</b> for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111... (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award <b>A0</b> for the final mark (and full <b>FT</b> is available in subsequent parts)

### 3 Implied marks

Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

### 4 Follow through marks (only applied after an error is made)

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then **FT** marks should be awarded for *their* correct answer, even when working is not present.

**For example:** following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is **(M1)A1**, it is possible to award full marks for *their* correct answer, **without working being seen**. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (e.g. probability greater than 1,  $\sin \theta = 1.5$ , non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word “their” in a description, to indicate that candidates may be using an incorrect value.
- If the candidate’s answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any **FT** marks in the subsequent parts. This includes when candidates fail to complete a “show that” question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these **FT** rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was “Hence”.

## 5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular misread. Use the **MR** stamp to indicate that this has been a misread and do not award the first mark, even if this is an **M** mark, but award all others as appropriate.

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. probability greater than 1,  $\sin \theta = 1.5$ , non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- **MR** can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

## 6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for parts of questions are indicated by **EITHER . . . OR**.

## 7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

## 8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come “*from the use of 3 sf values*”.

**Simplification of final answers:** Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed, and any

values that lead to integers should be simplified; for example,  $\sqrt{\frac{25}{4}}$  should be written as  $\frac{5}{2}$ . An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example,  $\frac{10}{4}$  may be left in this form or written as  $\frac{5}{2}$ .

However,  $\frac{10}{5}$  should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g.  $4e^{2x} \times e^{3x}$  should be simplified to  $4e^{5x}$ , and  $4e^{2x} \times e^{3x} - e^{4x} \times e^x$  should be simplified to  $3e^{5x}$ . Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so  $x(x+1)$  and  $x^2 + x$  are both acceptable.

**Please note:** intermediate **A** marks do NOT need to be simplified.

## 9 Calculators

A GDC is required for this paper, but if you see work that suggests a candidate has used any calculator not approved for IB DP examinations (eg CAS enabled devices), please follow the procedures for malpractice.

## 10. Presentation of candidate work

**Crossed out work:** If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

**More than one solution:** Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is “first”.

1. (a) (i) ( $Q_3 =$ ) 7.2 A2

**Note:** Award **A1A0** if the lower quartile, 6.5, is given as the answer.  
Award **A1A0** for a correct ordered list of happiness scores,  
when the correct  $Q_3$  is not seen.

(ii)  $Q_1 = 6.5$  (A1)  
 $IQR = 7.2 - 6.5$   
 $= 0.7$  A1

[4 marks]

(b)  $Q_3 + 1.5 \times IQR$  (A1)

$(7.2 + 1.5 \times 0.7 =) 8.25$  A1

since  $7.5 < 8.25$  R1

Switzerland is not an outlier AG

**Note:** Do not award **A0A0R1**.

[3 marks]

(c) (i)  $a = 3.5$  A1

(ii)  $b = 8$  A1

(iii)  $c = 3.5$  A1

[3 marks]

(d) (i) ( $r_s =$ ) 0.164 (0.164134...) A2

(ii) France rank (of sixth) is unchanged (so the  $r_s$  is unchanged) R1

[3 marks]

(e) Because  $r_s$  is too close to zero and hence Jose's conclusion is not appropriate R1

**Note:** Award **R1** for a comment on the value  $r_s$  and "not appropriate" as a conclusion.  
Accept " $r_s$  indicates a weak correlation of the ranks".  
Do not accept " $r_s$  indicates a weak correlation of happiness score and country rank".

[1 mark]

[Total 14 marks]



2. (a)  $\frac{1}{2} \times 1 \times ((4 + 0.7) + 2(7.3 + 6.7 + 4 + 1.3))$  (A1)(A1)

$= 21.7$  (21.65) (cm<sup>2</sup>) A1  
[3 marks]

(b) 22.1 (22.08333...,  $\frac{265}{12}$ ) (cm<sup>2</sup>) A2  
[2 marks]

(c)  $600 = \pi r^2 h$  A1  
[1 mark]

(d) **METHOD 1** (Substitution of  $h = \frac{600}{\pi r^2}$  in  $A = 2\pi r^2 + 2\pi r h$ )

$A = 2\pi r^2 + 2\pi r h$  (A1)

attempt to isolate  $h$  or  $\pi r h$  (M1)

$h = \frac{600}{\pi r^2}$  OR  $\pi r h = \frac{600}{r}$

correct substitution of their  $h$  into correct expression (M1)

$A = 2\pi r^2 + 2\pi r \left(\frac{600}{\pi r^2}\right)$   $\left(A = 2\pi r^2 + \frac{1200}{r}\right)$

$k = 1200$  A1

**METHOD 2** (Equating  $2\pi r h$  and  $\frac{k}{r}$ )

$A = 2\pi r^2 + 2\pi r h$  OR  $2\pi r h = \frac{k}{r}$  (A1)

attempt to isolate  $h$  or  $(\pi) r h$  or  $r$  or  $r^2$  (M1)

$h = \frac{600}{\pi r^2}$  OR  $\pi r h = \frac{600}{r}$  OR  $r = \sqrt{\frac{600}{\pi h}}$  OR  $r^2 = \frac{600}{\pi h}$

correct substitution of their  $h$  or  $\pi r h$  or  $r$  or  $r^2$  into  $2\pi r^2 + 2\pi r h = 2\pi r^2 + \frac{k}{r}$  (M1)

$2\pi r \left(\frac{600}{\pi r^2}\right) = \frac{k}{r}$  OR  $2 \left(\frac{600}{r}\right) = \frac{k}{r}$  OR  $2(600) = k$  OR  $2\pi \left(\frac{600}{\pi h}\right) h = k$

$k = 1200$  A1  
[4 marks]

continue...

Question 2 continued.

(e) (i)  $\frac{dA}{dr} = 4\pi r - 1200r^{-2}$

**A1(M1)A1**

**Note:** Award **A1** for  $4\pi r$  seen, and **(M1)** for expressing  $\frac{1200}{r}$  as  $1200r^{-1}$  (can be implied through  $\mp \frac{1200}{r^2}$  seen), **A1** for  $-1200r^{-2}$ . Award at most **A1(M1)A0** if any additional terms are seen.

(ii)  $0 = 4\pi r - 1200r^{-2}$  **OR**  $\frac{dA}{dr} = 0$

**(M1)**

$r = 4.57 \left( 4.570781\dots, \sqrt[3]{\frac{300}{\pi}} \right)$  (cm)

**A1**

**Note:** Award at most **M1A0** if the final answer is in terms of  $k$ .

**[5 marks]**

(f) ( $h =$ ) 9.14 (9.14156.....) (cm)

**A1**

( $C =$ )  $2\pi$  (4.570781.....)  $\approx 28.7$  (28.7190...)(cm)

**A1**

**EITHER**

the longest dimension of the label (9 cm) is less than both values and hence the label will fit (in any rotation)

**R1**

**OR**

$9 < 9.14$  and  $5 < 28.7$

**R1**

**Note:** Do not accept an argument based on the comparison of areas.

**[3 marks]**

**[Total 18 marks]**

3. (a) recognition of a geometric sequence (M1)

$$3\left(\frac{2}{3}\right)^4 \quad \text{OR} \quad 3, 2, 4/3 \dots$$

$$= 0.593 \text{ (0.592592\dots, } 16/27) \text{ (cm)}$$

**A1**  
**[2 marks]**

(b) attempt to sum a geometric sequence (M1)

$$\frac{3\left(1-\left(\frac{2}{3}\right)^5\right)}{1-\frac{2}{3}} \quad \text{OR} \quad 3+2+\frac{4}{3} \dots$$

$$= 7.81 \text{ (7.81481\dots, } 211/27) \text{ (cm)}$$

**A1**  
**[2 marks]**

(c) recognition of need to find sum to infinity (M1)

$$\frac{3}{1-\frac{2}{3}}$$

$$= 9 \text{ (cm)}$$

(A1)

**A1**  
**[3 marks]**

(d) Comparing the sum of the widths greater than (or equal to) 8.5 (M1)

$$\text{e.g. } \frac{3\left(1-\left(\frac{2}{3}\right)^n\right)}{1-\frac{2}{3}} \geq 8.5 \quad \text{OR} \quad \text{sketch} \quad \text{OR} \quad \text{list of values with cross-over values}$$

$$7.13 \text{ (7.1285338740543\dots) seen}$$

(A1)

$$n \geq 7.13$$

$$n = 8$$

**A1**  
**[3 marks]**

(e) attempt to divide two adjacent areas (M1)

$$\frac{4}{9} \text{ (0.444, 0.444444\dots)}$$

**A1**  
**[2 marks]**

*continue...*

Question 3 continued.

(f)  $u_1 = 13.5$  (may be seen in part (e))

(A1)

attempt to find the sum of their  $n$  terms, with their  $u_1$  and their  $r$

M1

$$S_8 = \frac{13.5 \left( 1 - \left( \frac{4}{9} \right)^8 \right)}{1 - \frac{4}{9}}$$

**Note:** Do not award **M1** if 5 or infinity used for  $n$  or if  $\frac{2}{3}$  used for  $r$ .

$$= 24.3 \text{ (24.2630...)} \text{ (cm}^2\text{)}$$

A1

**Note:** Accept 24.2 (24.2439...) from using 0.444.

[3 marks]

[Total: 15 marks]

4. (a) attempt to use Pythagoras' theorem (M1)

$$\sqrt{3.4^2 - 2^2}$$

$$= 2.75 \text{ (2.74954...)} \text{ (m)}$$

**A1**  
**[2 marks]**

(b) (i) **METHOD 1** (Use of  $\frac{1}{2} \times a \times b \times \sin(\theta)$ )

$$60^\circ \quad \text{(A1)}$$

attempt to find area of one triangle using  $\frac{1}{2} \times a \times b \times \sin(\theta)$  (M1)

$$\frac{1}{2} \times 2 \times 2 \times \sin(60^\circ)$$

$$\left( 6 \times \frac{1}{2} \times 2 \times 2 \times \sin(60^\circ) \right) = 10.4 \text{ (10.3923..., } 6\sqrt{3}) \text{ (m}^2\text{)} \quad \text{A1}$$

**Note:** Award **A0M0A0** for  $\frac{1}{2} \times 2 \times 2$  or equivalent.

**METHOD 2** (Use of altitude)

(altitude is)  $\sqrt{3}$  (A1)

attempt to find the area of one triangle using  $\frac{1}{2} \times b \times h$  with their altitude. (M1)

$$\frac{1}{2} \times 2 \times \sqrt{3}$$

$$\left( 6 \times \frac{1}{2} \times 2 \times \sqrt{3} \right) = 10.4 \text{ (10.3923..., } 6\sqrt{3}) \text{ (m}^2\text{)} \quad \text{A1}$$

**Note:** Award **A0M0A0** for  $\frac{1}{2} \times 2 \times 2$  or equivalent.

**METHOD 3** (Finding the area of a trapezoid)

(altitude of one trapezoid is)  $\sqrt{3}$  (A1)

attempt to find area of one trapezoid using  $\frac{1}{2} \times (a + b)h$  (M1)

$$\frac{1}{2} \times (2 + 4)\sqrt{3} \quad (3\sqrt{3})$$

$$\left( 2 \times \frac{1}{2} \times (2 + 4)\sqrt{3} \right) = 10.4 \text{ (10.3923..., } 6\sqrt{3}) \text{ (m}^2\text{)} \quad \text{A1}$$

*continue...*

Question 4 continued.

(ii)  $\frac{1}{3} \times 10.3923... \times 2.74954...$  (A1)  
 $= 9.52 \text{ m}^3$  (9.52470...) A1

**Note:** Units must be seen.

[5 marks]

(c)  $\cos(\widehat{MAT}) = \frac{2}{3.4}$  or correct equivalent (A1)  
 (MAT =)  $54.0^\circ$  (53.9681..., 0.941921... radians) A1

[2 marks]

(d) Angle YAX =  $180 - 53.9681... = 126.031...^\circ$  (A1)  
 Angle YXA =  $180 - 35 - 126.031... = 18.9681...^\circ$  (A1)

**Note:** These angles may be seen in the sine rule.

Attempt to substitute into sine rule (M1)

$$\frac{AY}{\sin(18.9681...)} = \frac{2.6}{\sin(126.031)}$$

AY = 1.05 (1.04503...) (m)

A1  
 [4 marks]

(e) **METHOD 1 COSINE RULE**  
 attempt to substitute into cosine rule to form a quadratic for YZ (M1)  
 $0.9^2 = YZ^2 + 1.04503...^2 - 2 \times 1.04503... \times YZ \times \cos(35)$  (A1)  
 YZ = 0.185 (0.184692...) (m), 1.53 (1.52739...) (m) A1A1

**Note:** Accept 0.191 (0.191313...) from use of 3 s.f. values.

**METHOD 2 SINE RULE**

attempt to substitute into sine rule to find  $\widehat{YZA}$

$$\frac{\sin(\widehat{YZA})}{1.04503...} = \frac{\sin(35^\circ)}{0.9}$$

$\widehat{YZA} = 41.7597... \text{ or } \widehat{YZA} = 138.240...$   
 $\widehat{ZAY} = 103.240... \text{ or } \widehat{ZAY} = 6.75972...$  (A1)

**Note:** Award A1 for any of these angles seen.

$$\frac{YZ}{\sin(\widehat{ZAY})} = \frac{0.9}{\sin(35^\circ)}$$

YZ = 0.185 (0.184692...) (m), 1.53 (1.52739...) (m)

A1A1  
 [4 marks]  
 [Total: 17 marks]

5. (a) recognition that speed is the magnitude of  $\begin{pmatrix} 50 \\ -33 \\ 0 \end{pmatrix}$  (M1)

$$\sqrt{50^2 + (-33)^2} \quad \text{OR} \quad \left| \begin{pmatrix} 50 \\ -33 \\ 0 \end{pmatrix} \right|$$

$$= 59.9 \text{ (km h}^{-1}\text{). (59.9082...)}$$

**A1**  
**[2 marks]**

- (b)  $\begin{pmatrix} 50 \\ -33 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} -15 \\ -20 \\ 0 \end{pmatrix} = -90$  (A1)

$$\cos \theta = \frac{-90}{59.9082... \times 25}$$
(M1)

$$\theta = 93.4^\circ \text{ (93.4450...}^\circ, 1.63092...)$$
**A1**  
**[3 marks]**

- (c)  $\vec{OD} = \begin{pmatrix} 200 \\ -100 \\ 0.02 \end{pmatrix} + t \begin{pmatrix} -15 \\ -20 \\ 0 \end{pmatrix}$  A1

**[1 mark]**

- (d) (i)  $\begin{pmatrix} 200 \\ -100 \\ 0.02 \end{pmatrix} + t \begin{pmatrix} -15 \\ -20 \\ 0 \end{pmatrix} = \begin{pmatrix} 152 \\ p \\ 0.02 \end{pmatrix}$  OR  $200 = -15t + 152$  (M1)

$$t_1 = 3.2 \left( \frac{16}{5} \right)$$
A1

- (ii)  $p = -164$  A1
- [3 marks]**

*continue...*

Question 5 continued.

(e) (i) attempt to find difference between the two position vectors

(M1)

$$\begin{pmatrix} 190 - 65t \\ -95 + 13t \\ 0.02 \end{pmatrix}$$

A1

**Note:** Award **A1M1A0** for  $\begin{pmatrix} -190 + 65t \\ 95 - 13t \\ -0.02 \end{pmatrix}$ .

(ii) attempt to find  $\left| \begin{pmatrix} 190 - 65t \\ -95 + 13t \\ 0.02 \end{pmatrix} \right|$

(M1)

$$\sqrt{(190 - 65t)^2 + (13t - 95)^2 + 0.02^2}$$

(A1)

attempt to find minimum. (e.g.  $t = 3.09$  hours)

(M1)

closest distance = 55.9 (55.8931...) (km)

A1

[6 marks]

[Total: 15 marks]



6. (a) anticlockwise rotation of  $15^\circ$  about the origin **A1**  
**[1 mark]**

(b) recognizing that  $l$  is equivalent to one rotation of  $360^\circ$  **(M1)**

e.g.  $\frac{360}{15}$   
 $= 24$

**A1**  
**[2 marks]**

(c) (i)  $(B =) \begin{pmatrix} 1.05 & 0 \\ 0 & 1.05 \end{pmatrix}$  **A1**

(ii)  $(B^{24} =) \begin{pmatrix} 3.23 & 0 \\ 0 & 3.23 \end{pmatrix}$  **(A1)**

**Note:** Award **A1** for 3.23 (3.22509...) **OR**  $1.05^{24}$ .

enlargement, with a scale factor of 3.23 ( $1.05^{24}$ ), (centre (0, 0))

**A1**  
**[3 marks]**

(d)  $C = \begin{pmatrix} 1.05 \cos(15^\circ) & -1.05 \sin(15^\circ) \\ 1.05 \sin(15^\circ) & 1.05 \cos(15^\circ) \end{pmatrix}$


$\left( = \begin{pmatrix} 1.01 \text{ (1.01422...)} & -0.272 \text{ (-0.271759...)} \\ 0.272 \text{ (0.271759...)} & 1.01 \text{ (1.01422...)} \end{pmatrix} = \frac{21}{80} \begin{pmatrix} \sqrt{6} + \sqrt{2} & -\sqrt{6} + \sqrt{2} \\ \sqrt{6} - \sqrt{2} & \sqrt{6} + \sqrt{2} \end{pmatrix} \right)$  **A2**

**Note:** Award **A1** for at least two correct elements in the matrix.

**[2 marks]**  
*continue...*

Question 6 continued.

(e)



anticlockwise spiral  
starting at (1, 0) and ending at (3.23, 0)

**A1**

**A1**

**Note:** Accept a continuous curve instead of discrete points. The two **A1** marks can be awarded independently

**[2 marks]**

(f)  $T \begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} p \\ q \end{pmatrix}$

**(M1)**

$0.9p + 2 = p$     **OR**     $0.8q + 1 = q$

**(A1)**

$p = 20, q = 5$     **OR**     $\begin{pmatrix} 20 \\ 5 \end{pmatrix}$

**A1**

**[3 marks]**

**[Total: 13 marks]**

7. (a)  $\frac{dy}{dt} = \frac{d^2x}{dt^2}$

**A1**

$$\frac{dy}{dt} + ay + bx = 0$$

**A1**

$$\frac{dy}{dt} = -bx - ay$$

**AG**

**[2 marks]**

(b)  $\begin{pmatrix} 0 & 1 \\ -77 & -18 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = -7 \begin{pmatrix} x \\ y \end{pmatrix}$  **OR**  $\begin{pmatrix} 0 & 1 \\ -77 & -18 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = -11 \begin{pmatrix} x \\ y \end{pmatrix}$

**(M1)**

$$y = -7x$$

eigenvector is  $\begin{pmatrix} 1 \\ -7 \end{pmatrix}$  (or any multiple)

**A1**

$$y = -11x$$

eigenvector is  $\begin{pmatrix} 1 \\ -11 \end{pmatrix}$  (or any multiple)

**A1**

**[3 marks]**

(c) (i)  $\begin{pmatrix} x \\ y \end{pmatrix} = Ae^{-7t} \begin{pmatrix} 1 \\ -7 \end{pmatrix} + Be^{-11t} \begin{pmatrix} 1 \\ -11 \end{pmatrix}$

**(A1)**

substitution of initial values

**(M1)**

two correct equations (not in vector form)

**(A1)**

$$5 = A + B, 2 = -7A - 11B$$

$$A = \frac{57}{4}, B = -\frac{37}{4}$$

**(A1)**


$$x = \frac{57}{4}e^{-7t} - \frac{37}{4}e^{-11t} \quad (x = 14.25e^{-7t} - 9.25e^{-11t})$$

**A1**

*continue...*

Question 7 continued.

(ii)



decreasing curve


A1

**Note:** There is a maximum at  $t = 0.00496$ , but this does not need to be shown.

starting at  $x=5$  and asymptote at  $x = 0$

A1  
[7 marks]

(d) (i)



spiral towards origin  
starting at (5, 2)

A1

A1

$$\frac{dy}{dt} = -85(5) - 18(2) = -461 < 0$$


Clockwise spiral

A1

continue...

Question 7 continued.

(ii)



oscillations  
starting at  $x = 5$ , decreasing amplitude

**A1**  
**A1**  
**[5 marks]**

(e) spring 1 is better as it gives a smoother ride (less oscillations, less bouncy)

**R1**  
**[1 mark]**  
**[Total:18 marks]**

---